



First report of a '*Candidatus* Phytoplasma solani'-related strain (16SrXII-A subgroup) associated with pistachio leaf scorch disease in Turkey

M.E. Guldur^{1*}, E. Simsek¹, B.K. Caglar², M. Dikilitas¹, H. Gumus¹ and H. Ayvaci¹

¹ Department of Plant Protection, Harran University, Sanliurfa, Turkey; ² Department of Plant Protection, Cukurova University, Adana, Turkey

*E-mail: mguldur@harran.edu.tr

Received: 18 Apr 2018. Published: 26 Jul 2018. Keywords: *Pistacia vera*, scorching, stolbur phytoplasma,

Pistachio (*Pistacia vera*) is a horticultural tree grown for its edible seeds. It is of great commercial importance for Turkey which is one of the main global producers. Pistachio trees are mostly cultivated in the Sanliurfa, Siirt, Gaziantep and Adiyaman provinces in Turkey. In recent years, symptoms regarding leaf scorch, chlorosis, and small leaves and significant yield loss have been observed in pistachio orchards.

Leaf samples were collected from symptom-bearing (Figs. 1-2) and symptomless pistachio trees in different pistachio orchards. Total nucleic acid was extracted using a CTAB-based method from each sample. Nucleic acids were tested with two-round PCR using P1/P7 primers in direct PCR followed by R16F2n/R2 primers in nested PCR to amplify the 16S rRNA gene (Duduk *et al.*, 2013). Nested-PCR amplicons of approximately 1.2 kb were obtained from all symptom-bearing samples but not from symptomless ones. Amplicons were purified, cloned and sequenced directly in both directions and sequences assembled and compared to reference strains using MEGA7 software (www.megasoftware.net). One representative isolate (2H2EF9) sequence was deposited in GenBank with the accession number MH088869. The isolate showed 99% identity to 16SrXII group phytoplasmas according to BLAST analysis. A computer-simulated RFLP analysis using 17 restriction enzymes with iPhyClassifier (Zhao *et al.*, 2009) and a phylogenetic tree generated by MEGA7 using the neighbour-joining method, confirmed the identification (Fig 3).

The association of phytoplasma (16SrI group) and pistachio was first reported by Marzachi *et al.* (1999) in Italy. There are also reports of phytoplasmas associated with distinct 16Sr groups infecting pistachio in Iran (Ghayeb Zamharir & Mirabolfathi, 2011), and Ghayeb Zamharir (2016; 2018) reported the presence of 16SrXII (associated with scorch disease) and 16SrIX groups of phytoplasma in Iran. Similarly, we detected and identified phytoplasmas in pistachio trees expressing leaf scorch symptoms in Sanliurfa, Turkey.

To the best of our knowledge, this is the first report of '*Ca. P. solani*'-related leaf scorch disease of pistachio in Turkey. Since the phytoplasmas belonging to subgroup 16SrXII-A have an extensive host

range, mixed farming in the region may, therefore, have a potential threat to pistachio. The distribution, epidemiology and impact of the disease on pistachio yield are currently under investigation.

Acknowledgements

The authors would like to thank HUBAK (Harran University Scientific Research Institution) for financial support.

References

- Duduk B, Paltrinieri S, Lee IM, Bertaccini A, 2013. Nested PCR and RFLP Analysis based on the 16S rRNA Gene. In: Dickinson M, Hodgetts J, eds. *Phytoplasma. Methods in Molecular Biology (Methods and Protocols)*, vol 938. Totowa, USA: Humana Press, 159-171. http://dx.doi.org/10.1007/978-1-62703-089-2_14
- Ghayeb Zamharir M, 2016. Association of a '*Candidatus* Phytoplasma solani'-related strain with pistachio in Iran. *New Disease Reports* **34**, 9. <http://dx.doi.org/10.5197/j.2044-0588.2016.034.009>
- Ghayeb Zamharir M, 2018. Molecular study of phytoplasmas associated with pistachio yellows in Iran. *Journal of Phytopathology*, 161-166. <http://dx.doi.org/10.1111/jph.12672>
- Ghayeb Zamharir M, Mirabolfathi M, 2011. Association of a phytoplasma with pistachio witches' broom disease in Iran. *Journal of Phytopathology* **159**, 60-62. <http://dx.doi.org/10.1111/j.1439-0434.2010.01697.x>
- Marzachi C, Alma A, d'Aquilio M, Minuto G, Boccardo G. 1999. Detection and identification of phytoplasmas infecting cultivated and wild plants in Liguria (Italian Riviera). *Journal of Plant Pathology* **81**, 127-136. <http://dx.doi.org/10.4454/jpp.v81i2.1057>
- Zhao Y, Wei W, Lee I-M, Shao J, Suo X, Davis RE, 2009. Construction of an interactive online phytoplasma classification tool, iPhyClassifier, and its application in analysis of the peach X-disease phytoplasma group (16SrIII). *International Journal of Systematic and Evolutionary Microbiology* **59**, 2582-2593. <http://dx.doi.org/10.1099/ijs.0.010249-0>



Figure 1



Figure 2




Figure 3

To cite this report: Guldur ME, Simsek E, Caglar BK, Dikilitas M, Gumus H, Ayvaci H, 2018. First report of a '*Candidatus* Phytoplasma solani'-related strain (16SrXII-A subgroup) associated with pistachio leaf scorch disease in Turkey. *New Disease Reports* **38**, 4. <http://dx.doi.org/10.5197/j.2044-0588.2018.038.004>

©2018 The Authors

This report was published on-line at www.ndrs.org.uk where high quality versions of the figures can be found.