



First record of Grapevine Pinot gris virus infecting *Vitis vinifera* in the United Kingdom

G. Silva^{1*}, J. Lecourt², G.R.G. Clover³ and S.E. Seal¹

¹ Natural Resources Institute, University of Greenwich, Chatham Maritime, Kent ME4 4TB, UK; ² NIAB EMR, New Road, East Malling, Kent ME19 6BJ, UK; ³ Royal Horticultural Society, RHS Garden Wisley, Woking, Surrey GU23 6QB, UK

*E-mail: g.silva@gre.ac.uk

Received: 01 Aug 2018. Published: 10 Aug 2018.

Grapevine Pinot gris virus (GPGV) is a member of the genus *Trichovirus*, and was first identified in grapevine (*Vitis vinifera*) cv. Pinot Gris in Italy in 2012 (Giampetruzzi *et al.*, 2012). Since then GPGV has been reported in several European countries as well as Australia, Canada, China, Korea and the USA (Bertazzon *et al.*, 2016). In April 2017, a survey of four geographically separated vineyards in the UK was done to investigate the presence of GPGV. A dormant cane was sampled at random from each of the four locations (Pinot Noir clones 119, 336, 792 and 924, reciprocally grafted upon Gravesac, SO4 or 3309 Couderc rootstocks).

RNA was extracted from cambial scrapings using a modified CTAB method (Abarshi *et al.*, 2010) and tested by RT-PCR using the specific primer pair Pg-Mer-F1 and Pg-Mer-R1 (Beuve *et al.*, 2015) targeting the movement protein (MP) gene. One sample, clone 336 grafted upon Gravesac, tested positive for GPGV yielding the expected 770 bp fragment. To confirm the detection of GPGV in the UK, a second RT-PCR was performed using GPGV-specific primers spanning the end of the MP and the beginning of the coat protein gene sequences (Morelli *et al.*, 2014). An amplicon of the expected size (588 bp) was obtained.

PCR amplicons from both GPGV primer sets were sequenced in both directions and the resulting sequences assembled using Geneious v10.2.6 (Biomatters Ltd., New Zealand). Sequences overlapped by 441 bp and generated a single contig of 917 bp representing a partial sequence of the GPGV isolate from the UK (GenBank Accession No. MG983746). Similarity searches using BLAST showed that this GPGV isolate shares the highest sequence identity at the nucleotide level (99%) with isolate GPGV Mer from France (KM491305).

To our knowledge, this is the first report of GPGV in the UK. GPGV has been associated with symptoms of leaf mottling and deformation in grapevine and is transmitted by the eriophyid mite *Colomerus vitis* (Malagnini *et al.*, 2016) which is present in the UK. Further large-scale studies should be done to determine the prevalence and spread of GPGV in the UK and evaluate the impact of the virus on yield and wine quality. The UK wine industry is a fast-growing sector with production projected to

increase from the current 6 million bottles of wine per annum to c. 40 million bottles by 2040. Extensive knowledge is needed about the presence and incidence of viruses in UK vineyards to develop efficient control strategies critical to enable the continued development of the industry.

References

- Abarshi MM, Mohammed IU, Wasswa P, Hillocks RJ, Holt J, Legg JP, Seal SE, Maruthi MN, 2010. Optimization of diagnostic RT-PCR protocols and sampling procedures for the reliable and cost-effective detection of *Cassava brown streak virus*. *Journal of Virological Methods* **163**, 353-359. <http://dx.doi.org/10.1016/j.jviromet.2009.10.023>
- Bertazzon N, Filippin L, Forte V, Angelini E, 2016. Grapevine Pinot gris virus seems to have recently been introduced to vineyards in Veneto, Italy. *Archives of Virology* **161**, 711-714. <http://dx.doi.org/10.1007/s00705-015-2718-2>
- Beuve M, Candresse T, Tannieres M, Lemaire O, 2015. First report of *Grapevine Pinot gris virus* (GPGV) in grapevine in France. *Plant Disease* **99**, 293. <http://dx.doi.org/10.1094/PDIS-10-14-1008-PDN>
- Giampetruzzi A, Roumi V, Roberto R, Malossini U, Yoshikawa N, La Notte P, Terlizzi F, Credi R, Saldarelli P, 2012. A new grapevine virus discovered by deep sequencing of virus- and viroid-derived small RNAs in cv Pinot gris. *Virus Research* **163**, 262-268. <http://dx.doi.org/10.1016/j.virusres.2011.10.010>
- Malagnini V, De Lillo E, Saldarelli P, Beber R, Duso C, Raiola A, Zanotelli L, Valenzano D, Giampietruzzi A, Morelli M, Ratti C, Causin R, Gualandri V, 2016. Transmission of grapevine Pinot gris virus by *Colomerus vitis* (Acari: Eriophyidae) to grapevine. *Archives of Virology* **161**, 2595-2599. <http://dx.doi.org/10.1007/s00705-016-2935-3>
- Morelli M, de Moraes Catarino A, Susca L, Saldarelli P, Gualandri V, Martelli GP, 2014. First report of Grapevine Pinot gris virus from table grapes in southern Italy. *Journal of Plant Pathology* **96**, 431-439. <http://dx.doi.org/10.4454/JPP.V96I2.039>

To cite this report: Silva G, Lecourt J, Clover GRG, Seal SE, 2018. First record of *Grapevine Pinot gris virus* infecting *Vitis vinifera* in the United Kingdom. *New Disease Reports* **38**, 7. <http://dx.doi.org/10.5197/j.2044-0588.2018.038.007>

©2018 The Authors

This report was published on-line at www.ndrs.org.uk where high quality versions of the figures can be found.